

CLINICAL MANUAL (IWK)

MATERNAL CHILD HEALTH MANUAL (NSHA)

Policy and Procedure

TITLE:	Skin-to-Skin Contact for Healthy Term Infants	NUMBER: IWK	IWK-1745
		NSHA	MC-NB-001
Sponsor:	Director Women’s and Newborn Health, Izaak Walton Killam (IWK) Health Centre Senior Director Maternal Child Health Nova Scotia Health Authority (NSHA)	Page:	1 of 11
Approved by:	IWK Medical Advisory Committee NSHA VP, Integrated Health Services, Primary Health Care Population Health	Approval Date IWK:	2018-01-04
		Effective Date IWK:	2018-01-04
		Approval Date NSHA:	2017-08-21
		Effective Date NSHA:	2018-05-22
Applies To:	IWK Staff and Health Care Providers NSHA Staff and Health Care Providers in Maternal/Newborn Health		

TABLE OF CONTENTS

PREAMBLE	2
POLICY STATEMENTS	2
GUIDING PRINCIPLES AND VALUES	3
PROCEDURE.....	3
REFERENCES	5
RELATED DOCUMENTS	7
Policies	7
Forms	7
Video	7
Brochures.....	7

Appendices	8
Replacing the Following District Health Authority Policies/Version History.....	8
Appendix A – Definitions	9
Appendix B – Initiation of Lactation: Anticipated Behaviours and Feeding Patterns*	10

PREAMBLE

1. IWK and NSHA Health Centre support evidence-based practice.
2. This policy promotes and supports mother and infant togetherness and Skin to Skin Contact (SSC) as recommended by:
 - The [Global Strategy for Infant and Young Child Feeding](#)
 - [UNICEF UK Baby-Friendly Initiative Standards](#)
 - [Breastfeeding Committee for Canada 10 Steps and WHO Code Outcome Indicators for Hospital and Community Services](#)
 - Accreditation Canada
 - The [Infant Feeding - Policy and Protocol - IWK - 1115 NSHA MC-GA-001](#)
 - The Operating Room Nurses Association of Canada (ORNAC) Standards, Guidelines, and Position Statements for Perioperative Registered Nurses

POLICY STATEMENTS

1. All Health Care Providers (HCPs), within the hospital and community supporting families during the prenatal, intrapartum and postpartum periods, must:
 - 1.1. Provide information to parents about the importance of immediate and uninterrupted SSC, encouraging discussion along the continuum of care.
 - 1.2. Promote and support the newborn being placed in SSC with their mother immediately following birth regardless of method of birth (vaginal or C-section) unless the mother or baby are medically unstable or safety is an issue.
 - 1.2.1. SSC must take place for at least an hour, until completion of the first feed, or as long as the mother wishes including on transfer to a different care area.
 - 1.2.2. Routine assessments of the newborn and mother must be completed without interrupting SSC.
 - 1.2.3. HCPs must document information about initial SSC on the permanent record. This includes information as to why SSC did not occur or was interrupted.
 - 1.3. Support Mother-Baby-Togetherness during the newborn period.

2. The newborn is removed only when the mother or baby are medically unstable and/or safety is an issue OR if requested by the mother based on her informed decision. SSC must begin/recommence as soon as medical status permits.
3. When the mother is unable to provide SSC secondary to medical issues or safety, the support person must be offered the opportunity to provide SSC until the mother is able and when safely possible; a person from the same home environment is optimal.

GUIDING PRINCIPLES AND VALUES

1. SSC maximizes physiological and psychological outcomes for both the mother and infant including optimal infant stability and infant attachment, as well as stimulation of feeding behavior.
 - 1.1. Optimal thermoregulation of the newborn is supported through uninterrupted SSC immediately following birth.
2. SSC is an effective non-pharmacological strategy for infant pain management.
 - 2.1. SSC **and** baby latched at the breast is the optimal non-pharmacological strategy for minor painful procedures in the neonatal period.
 - 2.2. All Staff and HCPs will center care on the dyad to support procedures that can be done safely in SSC.
3. The mother will be assessed for falls risk and education provided. (See relevant Fall Policies)

PROCEDURE

1. HCPs are responsible to:
 - 1.1. Support SSC immediately following birth by:
 - 1.1.1. Placing the newborn prone on the mother's chest with ventral surface of infant's body in direct contact with mother
 - 1.1.2. Drying infant's back and head (hat can be applied)
 - 1.1.3. Covering mother and baby with a warm, dry blanket while supporting practices that reduce the risk of infant overheating.
 - 1.2. Ensure safety of the newborn by
 - 1.2.1. Ensuring the newborn can spontaneously lift their head **at all times** to facilitate optimal breathing and attachment to breast
 - 1.2.2. Visually checking newborn's breathing and color when checking vital signs as per protocol
 - 1.2.3. Ensuring the newborn's nose and mouth are visible **at all times**
 - 1.2.4. Supporting SSC when mother drowsy/asleep by having a family member or nurse present to observe and ensure safety (falls prevention)

- 1.2.5. Asking the support person to hold the newborn's bottom or to firmly hold newborn's thigh to avoid slipping, when in the operating room
- 1.2.6. Raising mother's head of bed to 30 degrees or more following transfer
- 1.3. Obtain vital signs and observe the infant (temperature, breathing, color and tone) and mother as per standard practice in all care areas without interrupting SSC.
 - 1.3.1. Delay all routine measurements that necessitate the newborn being removed from SSC (such as weight and assessment for developmental dysplasia of the newborn's hips) until completion of the first feeding and ideally the completion of first hour of SSC.
 - 1.3.2. Delay routine medications until completion of the first feeding and ideally the completion of first hour of SSC.
 - 1.3.3. Newborn medications are administered with newborn in SSC and breastfeeding for any painful procedures to support pain management
- 1.4. Support SSC for painful procedures in addition to other supportive practices if the mother has decided to use human milk substitutes.
- 1.5. Use the approved IWK/NSHA Falls Prevention Tool ([See relevant Fall Policy](#)), to assess the mother's ability to safely provide SSC during mother's movement between stretcher, bed or wheelchair.
 - 1.5.1. If falls risk is low:
 - 1.5.1.1. The health care provider ensures the newborn is positioned vertically between mother's breasts; the mother crosses her arms around the newborn to hold baby securely.
 - 1.5.2. If falls risk is high:
 - 1.5.2.1. An alternate acceptable method is to place the newborn SSC on the support person, covered with a dry blanket, while the mother is transferred.
 - 1.5.2.2. As soon as the mother is settled, the support person re-replaces the infant SSC on the mother with the HCPs help.

NOTE: The health of the infant and mother is of utmost importance and if concern arises with the health of either, separation may be medically indicated.

The infant is removed only if medically indicated or requested by the mother and this is recorded in the permanent health care record.

SSC will begin/recommence as soon as medical status permits.

- 1.6. Be knowledgeable about the importance of SSC and provide education to families that promotes and supports SSC by:

- 1.6.1. Sharing approved resources and key messages (written and verbal) to families about SSC at all points of care. These messages include and are not limited to:
 - 1.6.1.1. The importance of immediate and uninterrupted SSC for at least the first hour, until completion of the first feed or as long as the mother desires ([See A Parent's Guide to Skin-to-Skin Contact Brochure](#))
 - 1.6.1.2. [9 Stages of the Newborn](#)
 - 1.6.1.3. Initiation of Lactation: Anticipated newborn behaviors and feeding patterns ([Appendix B](#))
 - 1.6.1.4. Infant cues and responsive parenting ([Loving Care Birth to 6 Months](#)),
 - 1.6.1.5. Positions that facilitates eye contact between mother and infant to support recognition of infant's cues
2. Managers are responsible to:
 - 2.1. Identify Staff and HCP training needs and provide support for education related to SSC, which includes expectations of Staff and HCPs in relation to the policy.
 - 2.2. Implement process to audit SSC practices immediately following birth (in different care team areas) at a minimum annually; Take corrective actions as needed. Data collected to include whether SSC is:
 - 2.2.1. Immediate and continuous for at least an hour, until completion of the first feed (at minimum)
 - 2.2.2. Delayed – include length of delay and medical reason why
 - 2.2.3. Discontinued – identify medical reason why
 - 2.2.4. Not at all (none) – identify medical reason why
3. IWK and NSHA Leadership are responsible to:
 - 3.1. Support development and implementation processes for the IWK/NSHA SSC Policy.

REFERENCES

- Accreditation Canada. Retrieved from <https://www.accreditation.ca>
- Anand KJS (2001). The international evidence-based group for neonatal pain. Consensus statement for the prevention and management of pain in the newborn. *Archives of Pediatrics and Adolescent Medicine*, 155 (2), 173-180.
- BC Women's Hospital and Health Centre. (2015). *Mother and baby togetherness position statement*. Retrieved from http://bccwhcms.medworxx.com/Site_Published/bcw/document_render.aspx?documentRender.IdType=29&documentRender.GenericField=1&documentRender.Id=16220
- Breastfeeding Committee for Canada. (2017). *The BFI 10 Steps and WHO Code Outcome Indicators for Hospitals and Community Health Services*. Retrieved from:

<http://breastfeedingcanada.ca/documents/Indicators%20-%20complete%20June%202017.pdf>

- Bystrova K (2008). Skin-to-skin contact and suckling in early postpartum: Effects on temperature, breastfeeding and mother-infant interaction. *A Study in St.Petersburg*, Doctoral thesis, Department for Women and Child Health, Karolinska Institutet, Stockholm, Sweden.
- Conde-Agudelo, A., Diaz-Rossello, J.L. (2014). Kangaroo mother care to reduce morbidity and mortality in low birth weight infants. *Cochrane Database Systematic Review*, Apr 22(4).
- CPS Position Statement. (Reaffirmed 2017). *The Baby-Friendly Initiative: Protecting, Promoting and Supporting Breastfeeding*. Retrieved from: <http://www.cps.ca/documents/position/baby-friendly-initiative-breastfeeding>
- Crenshaw J, Caldwell K, Brimdyr K, Widstrom AM, Svensson K, Champion J et al. (2012). Use of video-ethnographic intervention (PRECESS Immersion Method) to improve skin-to-skin care and breastfeeding rates. *Breastfeeding Medicine*, 7 (2), 68-79.
- Dani, C., Cecchi, A., Commare, A., Rapisardi, G., Breschi, R. & Pratesi, S. (2015). Behavior of the newborn during Skin-to-Skin. *Journal of Human Lactation*. January 2015.
- Dumas L, Lepage M, Bystrova K, Matthiesen AS, Welles-Nyström B, Widström AM. (2013). *Influence of SSC and rooming-in on early mother-infant interaction: a randomized controlled trial*. [Clinical Nursing Research](#), Aug 22(3), 310-36.
- Dumas L (2017). Ensuring skin-to-skin safety at vaginal and caesarean births. Presentation for staff at IWK Hospital, Halifax. September.
- Healthy Children Project. *The magical hour: holding your baby skin to skin in the first hour after birth*. 30 minute video and small book for parents. Available in both French and English at www.healthychildren.cc .
- Hedberg-Nyqvist K, Anderson CG, Bergman N, Cattaneo A, Charpak N, Davanzo R, Ewald U, Ibe O, Ludington-Hoe S, Mendoza S, Pallás-Allonso C, Ruiz-Peláez JG, Sizun J & Widström A-M (2010). Towards Universal Kangaroo Mother Care: recommendations and report from the First European conference and Seventh International Workshop on Kangaroo Mother Care. *Acta Paediatrica*, 99 (6), 820–826.
- Johnston, C., Campbell-Yeo, M., Fernandes, A., Inglis, D., Streiner, D. & Zee, R. (2010). Skin-to-Skin care for procedural pain in neonates (protocol). *Cochrane Database Systematic Review*, 31(1).
- Moore ER, Bergman N, Anderson GC & Medley N (2016). Early skin-to-skin contact for mothers and their healthy newborn infants. *Cochrane Database of Systematic Reviews*, 11 : CD003519.
- Operating Room Nurses Association of Canada. (2017). *The Operating Room Nurses Association of Canada (ORNAC) Standards, Guidelines, and Position Statements for Perioperative Registered Nurses* (13th Ed), Section 31(9), 334-342.

Province of Nova Scotia. (2015). [Breastfeeding Basics](#).

Province of Nova Scotia. (2014). [Loving Care: Birth to 6 months](#).

Province of Nova Scotia. (2013). *Family Friendly Breastfeeding Pledge*. Retrieved from: <http://novascotia.ca/dhw/healthy-development/documents/Family-Friendly-Breastfeeding-Pledge.pdf>

SOGC et al. Joint Policy Statement on Normal Childbirth, (2008). *The Journal of Obstetrics and Gynaecology Canada*, 30(12), 163–1165. Retrieved from: <https://sogc.org/wp-content/uploads/2013/01/gui221PS0812.pdf>

UNICEF. (2017). [Unicef UK Baby Friendly Initiative Standards](https://www.unicef.org.uk/babyfriendly/baby-friendly-resources/guidance-for-health-professionals/implementing-the-baby-friendly-standards/guide-to-the-baby-friendly-initiative-standards). Retrieved from <https://www.unicef.org.uk/babyfriendly/baby-friendly-resources/guidance-for-health-professionals/implementing-the-baby-friendly-standards/guide-to-the-baby-friendly-initiative-standards>

Widström A-M, Lilja G, Aaltomaa-Michalias P, Dahlöf A, Lintula M & Nissen E (2011). Newborn behaviour to locate the breast when skin-to-skin : A possible method for enabling early self-regulation. *Acta Paediatrica*, 100 (1), 79-85.

Widström AM, Ransjö-Arvidson AB & Christensson K (1993). *Breastfeeding is baby's choice*. Produced at Karolinska Institutet, Stockholm. Available in French, English and other languages via The Healthy Children Project.

World Health Organization. (2003). *Global Strategy for Infant and Young Child Feeding*. Retrieved from: <http://www.who.int/nutrition/publications/infantfeeding/9241562218/en/>

World Health Organization. (2003). *Kangaroo Mother Care A Practical Guide*. Retrieved from: <http://whqlibdoc.who.int/publications/2003/9241590351.pdf?ua=1>

RELATED DOCUMENTS

Policies

[IWK – 8665 – Kangaroo Care: SSC in the NICU](#)

[IWK – 1012 – Falls Prevention](#)

[NSHA CL-SM-010 Fall & Injury Prevention](#)

Forms

[Reproductive Care \(RCP\) Partogram](#)

Videos

The Healthy Children Project. (2010). Skin-to-Skin in the First Hour After Birth: Practical Advice for Staff after Vaginal and Cesarean Birth: Nine stages of the newborn.

[The Power of a Parent's Touch](#) (2014)

Brochures

[IWK/NSHA A Parent's Guide to SSC: What you can do to support your baby](#)

Appendices

[Appendix A](#) – Definitions

[Appendix B](#) – Initiation of Lactation: Anticipated Behaviours and Feeding Patterns

[Replacing the Following District Health Authority/IWK Policies/Version History](#)

* * *

Appendix A – Definitions

Health Care Providers	Care providers who provide direct patient care within hospital and communities who support families during the prenatal, intrapartum and postpartum periods.
Mother-Baby-Togetherness	Consists of keeping mother and infant in close proximity in the days following birth while encouraging SSC and combining mother-infant care. Mother-Baby-Togetherness results in improved attachment, communication, maternal self-confidence and emotional well-being.
Skin-to-Skin	When a mother holds her baby dressed only in a diaper on her bare chest, maximizing skin contact between the two. It may also be provided by a support person identified by the mother when extenuating circumstances prevent her from being able to provide it.
Staff	Includes all IWK and NSHA employees working within hospital and communities who support families during the prenatal, intrapartum and postpartum periods.
Support Person	Person identified by the mother.
Term Infant	Infant born at greater than or equal to (\geq) 37 completed weeks of gestation.

Appendix B – Initiation of Lactation: Anticipated Behaviours and Feeding Patterns*

Took Appendix 8.2 Initiation of Lactation: Anticipated Behaviours and Feeding Patterns from: BCC – *The BFI 10 Steps and WHO Code Outcome Indicators for Hospitals and Community Health Services*. July 19, 2017.

INFANT	Birth – 2 hours	2 – 20 hours	20 – 24 hours	24 – 48 hours	48 – 72 hours	Greater than 72 hours
State	Alert, eager to suckle	Periods of light and deep sleep	Increasing wakefulness followed by long, deep sleep after cluster feeds	Similar to 20 – 24 hours. Periods of deep and light sleep	Periods of light and deep sleep	Periods of deep and light sleep
Feeding patterns	Breastfeed within 1 – 2 hrs after birth but may feed minimally	Sporadic, variable and frequent feeds (offer Skin-to-Skin to maximize feeding opportunities)	Frequent or cluster feedings, which may occur during the night.	Feeds frequently (at least 8 times per day)	Feeds frequently (at least 8 times per day)	Feeds frequently (8 or more times a day)
Voids	Not usual	Increase as feedings increase. May void 0 – 1 times	Gradually increases, may void 0 – 1 times. (uric acid crystals can be normal)	Increasingly wet diapers, urine pale in color (may have 2 – 3 wet diapers in a day)	Increasingly wet diapers, urine pale (may have 3 or more wet diapers in a day)	Increasing numbers of wet diapers per day. Urine pale.
Stools	Not usual	Meconium	Meconium	Meconium and transition stool	Transition stools several times in the day	Transition – to lighter or yellowish stools
Weight		Decreases	Decreases	Decreases	Decreases	Decreases up to 10% and then begins to increase by day 4 or 5
MOTHER	Produces colostrum	Colostrum – as colostrum removed, alveoli cells secrete milk or colostrum	Colostrum – transition milk may start but this usually occurs earlier in multiples than primips	Transition milk – breast fullness may appear as milk starts to increase	Breast fullness	Engorgement if feedings have not been frequent

* Variances occur. Factors that slow initiation of the lactation process are: cesarean birth, analgesics and anaesthetics (epidurals included) during labour and birth, supplementation, lack of breast stimulation, sleepy infant, and any additional conditions that interfere with frequent and unlimited feedings. Roberta J. Hewat, PhD, RN, IBCLC May be copied in its entirety, not to be adapted without permission and not to be used for commercial purposes adapted with permission.

District Health Authority/IWK Policies Being Replaced

South West Health 100.138 Maintaining Parent-Infant Contact Following Cesarean Section

Version History

Major Revisions (e.g. Standard 4 year review)	Minor Revisions (e.g. spelling correction, wording changes, etc.)
New to NSHA & IWK 2018-02-28	TOC revised 2017-10-03
	2017-12-15 Added "Contact" to the title at the request of <i>Janine McClure RN BSc IBCLC</i>
	2018-01-16 Made the following changes based on feedback from IWK approval Process: <ul style="list-style-type: none"> • Spelled out (>) greater than in Appendix A and B • 1.2.1 'its' changed to 'their'
	2018-01-25made the following changes under "References": <ul style="list-style-type: none"> • Province of Nova Scotia. (2015). Breastfeeding Basics - inserted updated hyperlink and removed URL address • Province of Nova Scotia. (2014). Loving Care: Birth to 6 months - inserted updated hyperlink and removed URL address • UNICEF. (2013). Unicef UK Baby Friendly Initiative Standards - Date changed from 2013 to 2017
	2021-08-18 Policy statement number 2: <ul style="list-style-type: none"> • 5th bullet "Nova Scotia Provincial Breastfeeding Policy" was changed to "The Infant Feeding - Policy and Protocol - IWK - 1115 NSHA MC-GA-001"